

Breast Cancer: Addressing Disparities, Improving Care

Sarina Schrager, MD, MS, *WMJ* Associate Editor

Breast cancer remains the leading cause of new cancer diagnoses among women in the United States and the second leading cause of cancer deaths, behind lung cancer.¹ As with many other chronic diseases, there is a striking racial disparity in breast cancer mortality. And although overall mortality rates from breast cancer in all women have declined over the past 20 years—likely due to expanded screening programs and advances in treatment options—racial disparities continue to exist.

White women and black women have similar rates of new breast cancer diagnoses, yet black women have an approximately 30% higher chance of dying from breast cancer.^{1,2} Reasons for this disparity vary. Black women may have multiple barriers to care including lack of health insurance and poor access to mammography screening programs. In some cities, the quality of care may be erratic based on location. When diagnosed with breast cancer, black women are more likely to have a more advanced stage of disease as well as triple negative breast cancer (estrogen receptor, progesterone receptor, and HER2 receptor negative), which is harder to treat and carries a poorer prognosis.² All of these factors can impact survival.

Several interventions have been tried to decrease these disparities. Notably, the Metropolitan Chicago Breast Cancer Task Force, which was created in Chicago in 2008 used a comprehensive public health approach to reduce disparities in the community—previously among the highest in the country. The Task Force found a striking variability in qual-

ity of mammogram services in different parts of the city and higher uninsured rates for women of color. As an intervention, they provided free, high quality mammography, partnered with community groups to do education about breast cancer screening and ensured

and without insurance. The Community-Academic partnership employed a community advisory committee and used these community relationships to ensure buy-in from the community groups. In this way, the medical knowledge from the academic partner (Medical

...Although overall mortality rates from breast cancer in all women have declined over the past 20 years—likely due to expanded screening programs and advances in treatment options—racial disparities continue to exist.

that all women received quality care.³ This multipronged, public health approach reduced the disparity in breast cancer mortality between white and black women by close to half, bringing Chicago in line with data from the rest of the country.

The paper by Kamaraju, et al⁴ in this issue describes a similar intervention in Milwaukee. The authors partnered with community organizations, presented educational seminars about breast health, and provided information and transportation for women to get free mammograms through the Wisconsin Well Woman Program. They also enlisted a mobile mammography unit that enabled women to get mammography on site at the neighborhood community centers. Over a 2-year period, the project affected almost 500 women and documented significant increases in mammography rates in both women with

College of Wisconsin) was shared with patients in ways that they understood, in an environment where they felt comfortable. This study is a great example of an innovative method to bring health care to the community in order to improve care for underserved women.

Additionally in this issue, 2 studies focus on evidence-based care for women with breast cancer. The project reported by Hill, et al⁵ describes an intervention to ensure adherence to national guidelines for women diagnosed with early stage breast cancer. The new guidelines by the National Comprehensive Cancer Network, established in 2016, do not recommend doing screening lab tests in these women. Previously, all women diagnosed with early stage breast cancer routinely had complete blood cell count and liver profile measurements. Using a multipronged intervention that targeted providers and included educa-

tion, feedback, and positive reinforcement (gift cards), the authors successfully achieved over 80% compliance with the new guidelines. This paper is an example of an effective quality improvement initiative.

Teaching residents about quality improvement (QI) is the focus of the paper by Reardon et al.⁶ This paper describes the development of a successful QI curriculum for psychiatry residents. The curriculum involves a faculty development component, time to pursue projects, and linking QI projects to Maintenance of Certification within the discipline. Most of the residents who participated completed successful QI projects.

Finally, Chaudhary et al⁷ evaluate the predictive power of progesterone receptor status in recurrence rates among women with ductal carcinoma in situ (DCIS). Invasive breast cancers with positive estrogen receptor status but negative progesterone status are more aggressive than those with positive estrogen and progesterone receptor status. The authors evaluated whether that subtype of tumors (estrogen receptor positive and progesterone receptor negative) was predictive of increased rates of recurrence in women with DCIS. They followed a cohort of almost 700 women for 5 years and found that progesterone receptor status did not predict recurrence rates. This information can be helpful when counseling women with DCIS about treatment options.

Breast cancer is common, and women of color continue to experience poorer outcomes after being diagnosed. This issue of the journal summarizes important research from the public health and clinical care perspectives that can help bridge the disparity gap and lead to more equitable outcomes.

REFERENCES

1. US Cancer Statistics Working Group. US Cancer Statistics Data Visualizations Tool, based on November 2017 submission data (199-2015). Centers for Disease Control website. <https://gis.cdc.gov/Cancer/USCS/DataViz.html>. Published June 2018. Accessed June 11, 2018.
2. Yedjou CG, Tchounwou PB, Payton M, et al. Assessing the racial and ethnic disparities in breast cancer mortality in the United States. *Int J Environ Res Public Health*. 2017;14(5). doi:10.3390/ijerph14050486.
3. Sighoko D, Murphy AM, Irizarry B, Rauscher G, Ferrans C, Ansell D. Changes in the racial disparity in breast cancer mortality in the ten US cities with the largest African American populations from 1999 to 2013: the reduction in breast cancer mortality disparity in Chicago. *Cancer Causes Control*. 2017;28(6):563-568.
4. Kamarju S, DeNomie M, Visotcky A, et al. Increasing mammography uptake through academic-community partnerships targeting immigrant and refugee communities in Milwaukee. *WMJ*. 2018;117(2):55-61.
5. Hill LA, Vang CA, Kennedy CR, et al. A strategy for changing adherence to national guidelines for decreasing laboratory testing for early breast cancer patients. *WMJ*. 2018;117(2):68-72.
6. Reardon CL, Creado S, Hafer R, et al. A curriculum for residents to develop successful quality improvement projects. *WMJ*. 2018;117(2):79-82.
7. Chaudhary LN, Jawa Z, Hanif A, et al. Does progesterone receptor matter in the risk of recurrence for patients with ductal carcinoma in situ? *WMJ*. 2018;117(2):62-67.

WMJ

Let us hear from you!

If an article strikes a chord or you have something on your mind related to medicine, we want to hear from you. Submit your letter via e-mail to wmj@wismed.org or send it to:

WMJ Letters • 330 E Lakeside St • Madison, WI 53715

VA



U.S. Department
of Veterans Affairs
Veterans Health
Administration

Minneapolis VA Health Care System BE/BC Family Practice or Internal Medicine

Chippewa Falls, WI
Superior, WI

Our Outpatient Clinics are open Monday through Friday, 8 a.m. to 4:30 p.m., closed on all federal holidays. No weekend duties and no call. Assigned panel of patients, panel size is determined by VHA PCMM software package, panel size is specific to their practice and setting. No emergency services on site.

Must have a valid medical license anywhere in the US. Background check required. BC and 2 to 3 years clinical experience with current BLS certification preferred.

VA providers enjoy an excellent benefits package and a state-of-the-art electronic medical record. Predictable work hours and conditions, VA physicians are able to take time for themselves and their families.

*Competitive salary
Paid malpractice insurance
Recruitment incentive possible
Annual Performance Pay bonus*

For more information:
Rick Pope
CBOC Business Manager
Richard.Pope@va.gov
612-467-1264
EEO employer

advancing the art & science of medicine in the midwest

WMJ

WMJ (ISSN 1098-1861) is published through a collaboration between The Medical College of Wisconsin and The University of Wisconsin School of Medicine and Public Health. The mission of *WMJ* is to provide an opportunity to publish original research, case reports, review articles, and essays about current medical and public health issues.

© 2018 Board of Regents of the University of Wisconsin System and The Medical College of Wisconsin, Inc.

Visit www.wmjonline.org to learn more.